

PART 1: QUESTIONS

Name: _____ Age: _____ Id: _____ Course: _____

Basic Concepts - Exam 2**Lesson: 4-6****Instructions:**

- Please begin by printing your Name, your Age, your Student Id , and your Course Name in the box above and in the box on the solution sheet.
- You have 90 minutes (class period) for this exam.
- You can not use any calculator, computer, cellphone, or other assistance device on this exam. However, you can set our flag to ask permission to consult your own one two-sided-sheet notes at any point during the exam (You can write concepts, formulas, properties, and procedures, but questions and their solutions from books or previous exams are not allowed in your notes).
- Each multiple-choice question is worth 5 points and each extra essay-question is worth from 0 to 5 points. (Even a simple related formula can worth some points).
- Set up your flag if you have a question.
- Relax and use strategies to improve your performance.

Exam Strategies to get the best performance:

- Spend 5 minutes reading your exam. Use this time to classify each Question in (E) Easy, (M) Medium, and (D) Difficult.
- Be confident by solving the easy questions first then the medium questions.
- Be sure to check each solution. In average, you only need 30 seconds to test it. (Use good sense).
- Don't waste too much time on a question even if you know how to solve it. Instead, skip the question and put a circle around the problem number to work on it later. In average, the easy and medium questions take up half of the exam time.
- Solving the all of the easy and medium question will already guarantee a minimum grade. Now, you are much more confident and motivated to solve the difficult or skipped questions.
- Be patient and try not to leave the exam early. Use the remaining time to double check your solutions.

1. Solve:

$$3(x - 2) - (x - 3) = -1$$

- a) 1
- b) 2
- c) 3
- d) 4
- e) There is no solution.

2. Solve:

$$(2x - 1) - 3(3 - x) = 10$$

- a) 1
- b) 2
- c) 3
- d) 4
- e) There is no solution.

3. Linda wants to walk dogs for \$15 an hour and work 5 hours per day. A total of 40% of her salary is for her own personal expenses. How many days must she work to save up \$450 per month?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 40 days
- e) None of the above.

4. An exam of 40 questions has the following rules:

- 3 points for each correct answer.
- -1 point for each incorrect answer.

How many correct answers do you need to get 40 points on the exam?

- a) 20 correct answers.
- b) 30 correct answers.
- c) 40 correct answers.
- d) 42 correct answers.
- e) None of the above.

5. Solve:

$$\frac{x-2}{x+1} - \frac{x-3}{x-1} = \frac{3}{x^2-1}; x \in \mathbb{R}$$

- a) 0
- b) 1
- c) 2
- d) 3
- e) There is no solution.

6. Solve:

$$\frac{x}{x-2} + 2x = \frac{2x^2-6}{x-2}; x \in \mathbb{R}$$

- a) There are infinite solutions.
- b) There are only one solution
- c) There is no solution..
- d) There are two solutions.
- e) There are three solutions.

7. Let x_1 and x_2 be the solutions of the equation, $ax^2 + bx + c = 0$; $a \neq 0$.

I. $x_1 \cdot x_2 = \frac{b}{a}$

II. $x_1 \cdot x_2 = \frac{c}{a}$

III. $ax^2 + bx + c = a(x + x_1)(x + x_2)$

Then:

- a) Only I is true.
- b) Only II is true.
- c) Only III is true.
- d) Only I and II are true.
- e) I, II, and III are true.

8. $ax^2 + bx + c = 0$ where a , b , and c are real constants with $a \neq 0$ and x is a variable such that $x \in \mathbb{R}$.

- I. This equation could have no solution.
- II. This equation could have three distinct solutions.
- III. This equation could have two distinct solutions.
- IV. This equation could have one distinct solution.

- a) I is false.
- b) II is false.
- c) III is false.
- d) IV is false.
- e) None of the above.

9. The solutions of the equation, $x^2 - 11x + 30 = 0$ are:

- a) $x_1 = 1$ and $x_2 = 2$
- b) $x_1 = 2$ and $x_2 = 3$
- c) $x_1 = 3$ and $x_2 = 4$
- d) $x_1 = 4$ and $x_2 = 5$
- e) None of the above.

10. Which equation below has solutions $x_1 = -6$ and $x_2 = 2$

- a) $x^2 + 4x + 12 = 0$
- b) $x^2 - 4x + 12 = 0$
- c) $x^2 + 2x - 12 = 0$
- d) $x^2 - 2x - 12 = 0$
- e) None of the above.

11. The solutions of $x^2 - 2x - 10 = 0$ are

- a) $x_1 = 1 - \sqrt{2}$ and $x_2 = 1 + \sqrt{2}$
- b) $x_1 = 1 - \sqrt{5}$ and $x_2 = 1 + \sqrt{5}$
- c) $x_1 = 1 - \sqrt{7}$ and $x_2 = 1 + \sqrt{7}$
- d) $x_1 = 1 - \sqrt{11}$ and $x_2 = 1 + \sqrt{11}$
- e) None of the above.

12. Given $x^2 - x + 4 = 0$, the Sum (S) and the Product (P) of the solutions are

- a) $S = 4$ and $P = 1$
- b) $S = 1$ and $P = 4$
- c) $S = 4$ and $P = -1$
- d) $S = -4$ and $P = -1$
- e) None of the above.

13. Let x_1, x_2 , and x_3 be the solutions of the equation $x^3 - 2x^2 + x = 0$. Then:

- a) $x_1 + x_2 + x_3 = 0$
- b) $x_1 + x_2 + x_3 = 1$
- c) $x_1 + x_2 + x_3 = 2$
- d) $x_1 + x_2 + x_3 = 3$
- e) $x_1 + x_2 + x_3 = 4$

14. Given the equation $\sqrt{4x+9} = x-3$, $x \in \mathbb{R}$. The sum of all possible solutions is:

- a) 0
- b) 8
- c) 10
- d) 12
- e) None of the above.

15. Let x_1 and x_2 be the solutions to $x^2 + 2x - 4 = 0$. Consider the equation:

- I. $x_1 + x_2 = 2$
- II. $x_1 \cdot x_2 = 4$
- III. $x_1^2 + x_2^2 = 12$
- IV. $x_1^3 + x_2^3 = -30$

Then

- a) Only I is correct.
- b) Only II is correct.
- c) Only III is correct.
- d) Only IV is correct.
- e) None of the above.

16. Solve: $5x + 3 > x + 2$. The solution is:

- a) $S = \{x \in \mathbb{R} / x > -2\}$
- b) $S = \{x \in \mathbb{R} / x > -1\}$
- c) $S = \{x \in \mathbb{R} / x > 0\}$
- d) $S = \{x \in \mathbb{R} / x > -2\}$
- e) None of the above.

17. Solve: $\frac{x-3}{x+1} \geq 0$. The solution is:

- a) $S = \{x \in \mathbb{R} / x > -1\}$
- b) $S = \{x \in \mathbb{R} / x < -1 \text{ or } x \geq 3\}$
- c) $S = \{x \in \mathbb{R} / x < -3 \text{ or } x > 3\}$
- d) $S = \{x \in \mathbb{R} / x < -3 \text{ or } x > 1\}$
- e) None of the above.

18. Solve: $\frac{x^2+x-6}{x} > 0$. The solution is:

- a) $S = \{x \in \mathbb{R} / -3 < x < 0 \text{ or } x > 2\}$
- b) $S = \{x \in \mathbb{R} / x < -3 \text{ or } 0 < x < 2\}$
- c) $S = \{x \in \mathbb{R} / x \leq -3 \text{ or } 0 \leq x \leq 2\}$
- d) $S = \{x \in \mathbb{R} / -3 \leq x < 0 \text{ or } x \geq 2\}$
- e) None of the above.

19. Solve: $-x^2 - x - 1 < 0$. The solution is:

- a) $S = \emptyset$ (There is no solution)
- b) $S = \mathbb{R}$
- c) $S = \{x \in \mathbb{R} / x > 0\}$
- d) $S = \{x \in \mathbb{R} / x < 0\}$
- e) None of the above.

20. Solve: $\frac{x(5-x)}{(5-x)} \geq 0$. The solution is:

- a) $S = \{x \in \mathbb{R} / x > 0 \text{ and } x \neq 5\}$
- b) $S = \{x \in \mathbb{R} / x \geq 0 \text{ or } x \neq -5\}$
- c) $S = \{x \in \mathbb{R} / 0 \leq x < 5 \text{ and } x > 5\}$
- d) $S = \{x \in \mathbb{R} / 0 < x < 5 \text{ and } x > 5\}$
- e) None of the above.

PART 2: SOLUTIONS

Consulting

Name: _____ Age: _____ Id: _____ Course: _____

Multiple-Choice Answers

Questions	A	B	C	D	E
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
11					
12					
13					
14					
15					
16					
17					
18					
19					
20					

Extra Questions

21. Solve: $x^4 - 8x^2 - 9 = 0$; $x \in \mathbb{R}$.

22. Solve: $\frac{1}{x} + x \geq -2$; $x \in \mathbb{R}$.

Let this section in blank

	Points	Max
Multiple Choice		100
Extra Points		25
Consulting		10
Age Points		25
Total Performance		160
Grade		A

23. Let x_1 and x_2 be the solution of the equation $x^2 - 2x - 1 = 0$.

$$\text{Calculate } \frac{x_1}{x_2} + \frac{x_2}{x_1} = \frac{x_1^2 + x_2^2}{x_1 x_2}.$$

24. Solve: $\frac{2-x}{4} = \frac{x-1}{5} + 1 ; x \in \mathbb{R}$.

25. Given:

1) $X \star Y = XY$, where $X, Y \in \mathbb{N}$.

Example: $2 \star 3 = 23$

2) $X \odot Y = X \star Y + X + Y$, where $X, Y \in \mathbb{N}$.

Example: $2 \odot 3 = 2 \star 3 + 2 + 3$

$$= 23 + 5$$

$$= 28$$

Find: $4 \odot 7 = ?$